

The Dependable Choice For **Acidity & Ulcer Pain**

# Oxysil



<b>Oxysil S</b>	Sucralfate U.S.P. 1000mg + Oxetacaine B.P. 20mg	<b>Syrup</b>
<b>Oxysil Forte</b>	Sucralfate USP 1gm Sugar Free	<b>Syrup</b>
<b>Oxysil Gel</b>	Aluminium Hydroxide I.P. 0.291gm + Magnesium Hydroxide I.P. 98mg + Oxetacaine B.P. 10mg	<b>Syrup</b>

### Sucralfate

- ✓ Useful in the treatment of gastric and duodenal ulcer, it is cytoprotective and able to increase the bioavailability of several growth factors, modulating the wound healing process.
- ✓ Reacts with gastric acid to form a long acting acid buffer for as long as 6 to 8 hours after a single dose.
- ✓ It also attaches to proteins on the surface of ulcer forming protective barriers at the ulcer surface, preventing further damage from acid pepsin & bile.

### Oxetacaine

- ✓ Is a potent local anesthetic that is administered orally for the relief of pain associated with peptic ulcer disease or esophagitis.
- ✓ Is highly active even in highly acidic conditions.



<b>GERD</b>	<b>Duodenal Ulcer</b>
<b>Stress Ulcer</b>	<b>Reflux Oesohagitis</b>